

Subject Index (2007-2008)

Editorial

Editorial.

Sekimizu K

2007; 1(1):1.

News

China-Japan enhance joint research cooperation for drug discoveries and development: News from CJMWDDT 2007 in Jinan, China.

Liu XY, Qu XJ, Tang W

2007; 1(1):2.

China's new drug R&D is steadily advancing.

Xu WF

2007; 1(1):3.

China's efforts to shed its title of "Leader in liver disease".

Li X, Xu WF

2007; 1(2):84-85.

Tes, a potential Mena-related cancer therapy target.

Li X

2008; 2(1):1.

Japan-China Joint Medical Workshop on Drug Discoveries and Therapeutics 2008: The need of Asian pharmaceutical researchers' cooperation.

Nakata M, Tang W

2008; 2(5):262-263.

Policy Forums

Dynamic of modernizing traditional Chinese medicine and the standards system for its development.

Gai RY, Xu HL, Qu XJ, Wang FS, Lou HX, Han JX, Nakata M, Kokudo N, Sugawara Y, Kuroiwa C, Tang W

2008; 2(1):2-4.

Three decades of GMP implementation in Thailand: Hardships and success.

Chanprapaph S

2008; 2(1):5-6.

Reviews

Viral infectious disease and natural products with antiviral activity.

Kitazato K, Wang Y, Kobayashi N

2007; 1(1):14-22.

Target validation: A door to drug discovery.

Chen XP, Du GH

2007; 1(1):23-29.

Therapeutic potential of heat-processed *Panax ginseng* with respect to oxidative tissue damage.
Yokozawa T, Kang KS, Yamabe N, Kim HY
2007; 1(1):30-44.

Secondary metabolites from higher fungi in China and their biological activity.
Liu JK
2007; 1(2):94-103.

Progress in cell membrane chromatography.
He LC, Wang SC, Yang GD, Zhang YM, Wang CH, Yuan BX, Hou XF
2007; 1(2):104-107.

Derivatization reagents in liquid chromatography/electrospray ionization tandem mass spectrometry for biomedical analysis.
SantaT, Al-Dirbashi OY, Fukushima T
2007; 1(2):108-118.

Sensing and reacting to dangers by caspases: Caspase activation via inflammasomes.
Takeishi A, Kuranaga E, Miura M
2008; 2(1):14-23.

Roles of the Duffy antigen and glycophorin A in malaria infectionand erythrocyte.
Hamamoto H, Akimitsu N, Arimitsu N, Sekimizu K
2008; 2(2):58-63.

Dermal drug delivery: Revisited.
Khandavilli S, Panchagnula R
2008; 2(2):64-73.

S-II mediated gene regulation.
Tano K, Nagata M, Akimitsu N
2008; 2(3):136-139.

Microfabrication-derived DDS: From batch to individual production.
Takada K
2008; 2(3): 140-155.

Spermicidal agents.
Shah HC, Tatke P, Singh KK
2008; 2(4):200-210.

Pulmonary drug delivery: Implication for new strategy for pharmacotherapy for neurodegenerative disorders.
Islam N, Rahman S
2008; 2(5):264-276.

Role of multidrug resistance associated proteins in drug development.
Zhou SF
2008; 2(6):305-332.

Pharmacogenomics-based clinical studies using a novel, fully automated genotyping system.
Hasegawa S, Kimura S, Kobayashi A, Imanishi N
2008; 2(6):333-338.

Brief Reports

Analysis on productivity of clinical studies across — Asian countries a case comparison.

Takahashi K, Sengoku S, Kimura H

2007; 1(1):4-8.

Lack of polymorphisms in the coding region of the highly conserved gene encoding transcription elongationfactor S-II (TCEA1).

Ito T, Doi K, Matsumoto N, Kakihara F, Noiri E, Hasegawa S, Tokunaga K, Sekimizu K

2007; 1(1):9-11.

GMP implementation in China: A double-edged sword for the pharmaceutical industry.

Gai RY, Qu XJ, Lou HX, Han JX, Cui SX, Nakata M, Kokudo N, Sugawara Y, Kuroiwa C, Tang W

2007; 1(1):12-13.

Study of drug resistance among 78 antiretroviral treatment-naïve patients with HIV-1 subtype B infection in central China.

Wang JQ, Huang XJ, Zhang HW, Li ZC, Wang XC, Li HY, Shen CL, Guo CP, Wei FL, Zhang T, Wu H

2007; 1(2):86-88.

A silkworm model of pathogenic bacterial infection.

Kaito C, Sekimizu K

2007; 1(2):89-93.

Establishment of a new cell line for performing sensitive screening of nuclear export inhibitors.

Watanabe K, Noda S, Kobayashi N

2008; 2(1):7-9.

Localization of glucagon-like peptide-2 receptor mRNA expression at different sites in the small intestine of rats.

Ikeda K, Myotoku M, Kurokawa N, Hirotani Y

2008; 2(1):10-13.

On the temperature dependence of the unbound drug fraction in plasma: Ultrafiltration method may considerably underestimate the true value for highly bound drugs.

Berezhkovskiy LM

2008; 2(2):74-76.

Novel N-hydroxybenzamide histone deacetylase inhibitors as potential anti-cancer agents.

Jiao J, Fang H, Xu WF

2008; 2(4):211-215.

Synthesis and antifungal activity of 3-substituted thiochromanones.

Liu Y, Luo W, Sun L, Guo C

2008; 2(4):216-218.

Effect of drug-polymer binary mixtures on the *in-vitro* release of ibuprofen from transdermal drug-in-adhesive layers.

Ho KY, Ord M, Dodou K

2008; 2(5):277-281.

Effect of benzyl-N-acetyl- α -galactosaminide on KL-6 mucin expression and invasive properties of a human pancreatic carcinoma cell line.

Xu HL, Inagaki Y, Wang FS, Kokudo N, Nakata M, Tang W

2008; 2(5):282-285.

Antioxidant activity of wild plants collected in Beni-Sueif governorate, Upper Egypt.

Abouzid S, Elshahaat A, Ali S, Choudhary MI
2008; 2(5):286-288.

Apoptosis-inducing effect of cinobufacini, *Bufo bufo gargarizans* Cantor skin extract, on human hepatoma cell line BEL-7402.

Qi FH, Li AY, Lv H, Zhao L, Li JJ, Gao B, Tang W
2008; 2(6):339-343.

Original Articles

Anti-depressant and anti-nociceptive effects of 1,4-benzodiazepine-2-ones based cholecystokinin (CCK₂) antagonists.

Lattmann E, Sattayasai J, Lattmann P, Billington DC, Schwalbe CH, Boonprakob J, Airarat W, Singh H, Offel M, Staaf A
2007; 1(1):45-56.

ONIOM DFT/PM3 calculations on the interaction between dapivirine and HIV-1 reverse transcriptase, a theoretical study.

Liang YH, Chen FE
2007; 1(1):57-60.

Studies on the development of rapidly disintegrating hyoscine butylbromide tablets.

Khattab IS, Zaghloul AA, Afouna MI
2007; 1(1):61-64.

Investigation of the binding behaviors of isonucleoside-incorporated oligonucleotides with complementary sequences.

Guan Z, Jin HW, Yang ZJ, Zhang LR, Zhang LH.
2007; 1(1):65-72.

Modification of 15-akylidene andrographolide derivatives as alpha-glucosidase inhibitor.

Xu HW, Liu GZ, Dai GF, Wu CL, Liu HM
2007; 1(1):73-77.

Evaluation of transdermal permeability of pentoxyfylline gel: *in vitro* skin permeation and *in vivo* microdialysis using Wistar rats.

Yan KS, Yan TX, Guo H, Li JZ, Wei LL, Wang C, Nie SF, Pan WS.
2007; 1(1):78-83.

Synthesis and cytotoxic activity of 3-phenyl-2-thio-quinoxaline 1,4-dioxide derivatives in hypoxia and in normoxia.

Sheng R, Xu Y, Weng QJ, Xia Q, He QJ, Yang B, Hu YZ
2007; 1(2):119-123.

A novel method to suppress the dispersal of Japanese cedar pollen by inducing morphologic changes with weak alkaline solutions.

Ishii K, Hamamoto H, Sekimizu K
2007; 1(2):124-129.

The effects of antidepressant drug on ethanol-induced cell death.

Johnson S, Williams AN, Johnson C, Ou XM
2007; 1(2):130-135.

A simple and rapid high performance liquid chromatography method to determine levofloxacin in human plasma and its use in a bioequivalence study.

Gao XX, Yao GC, Guo N, An F, Guo XJ
2007; 1(2):136-140.

Pharmacologic action of oseltamivir on the nervous system.

Ishii K, Hamamoto H, Sasaki T, Ikegaya Y, Yamatsugu K, Kanai M, Shibasaki M, Sekimizu K
2008; 2(1):24-34.

Immunosuppressive effect of ER-38925, a retinoic acid receptor subtype α -selective agonist, in mouse models of human graft-vs-host disease.

Hida T, Shikata K, Tokuhara N, Ishibashi A, Nagai M, Yamauchi T, Kobayashi S
2008; 2(1):35-44.

A new method of preparing TRH derivative-loaded poly(dl-lactide-coglycolide) microspheres based on a solid solution system.

Matsumoto A, Matsukawa Y, Nishioka Y, Harada M, Horikiri Y, Yamahara H
2008; 2(1):45-51.

3D-QSAR study with pharmacophore-based molecular alignment of hydroxamic acid-related phosphinates that are aminopeptidase N inhibitors.

Zhu HW, Fang H, Wang L, Hu WX, Xu WF
2008; 2(1):52-57.

Hsc70 regulates the nuclear export but not the import of influenza viral RNP: A possible target for the development of anti-influenza virus drugs.

Watanabe K, Takizawa N, Noda S, Tsukahara F, Maru Y, Kobayashi N
2008; 2(2):77-84.

UVB-dependent generation of reactive oxygen species by catalase and IgG under UVB light: Inhibition by antioxidants and anti-inflammatory drugs.

Murakami M, Taniguchi M, Takama M, Cui JH, Oyanagui Y
2008; 2(2):85-93.

Formulation and hypoglycemic activity of pioglitazone-cyclodextrin inclusion complexes.

Elbary AA, Kassem MA, Abou Samra MM, Khalil RM
2008; 2(2):94-107.

Reconstituted powder for suspension of antitubercular drugs formulated as microspheres for pediatric use.

Samad A, Sultana Y, Khar RK, Aqil M, Chuttani K, Mishra AK
2008; 2(2):108-114.

Synthesis and biological evaluation of substituted phenylpyrazole[4,5-*b*]oleanane derivatives as inhibitors of glycogen phosphorylase.

Chen J, Gong YC, Liu J, Zhang LY, Hua WY, Sun HB
2008; 2(2):115-121.

Using factorial design to improve the solubility and *in-vitro* dissolution of nimesulide hydrophilic polymer binary systems.

Khattab IS, Al-Saidan SM, Nada AH, Zaghloul AA
2008; 2(2):122-127.

Iontophoretic delivery of 5-fluorouracil through excised human stratum corneum.

Singh BN, Jayaswal SB
2008; 2(2):128-135.

Cholecystokinin antagonists (part 1): Antinociceptive, anxiolytic and antidepressant effects of *N*-(5-methyl-3-oxo-1,2-diphenyl-2,3-dihydro-1*H*-pyrazol-4-yl)-*N'*-phenylureas and carboxamides.

Lattmann E, Sattayasai J, Boonprakob Y, Singh H, Lattmann P, Dunn S
2008; 2(3):156-167.

Injection of cell-wall skeleton of *Mycobacterium bovis* BCG draining to a sentinel lymph node eliminates both lymph node metastases and the primary transplanted tumor.

Kashiwazaki Y, Murata M, Sato T, Miyauchi M, Nakagawa M, Fukushima A, Chiba N, Azuma I, Yamaoka T
2008; 2(3):168-177.

Immune response against cell-wall skeleton of *Mycobacterium bovis* BCG at the inoculation site and peripheral lymphoid organs.

Kashiwazaki Y, Murata M, Fujii T, Nakagawa M, Fukushima A, Chiba N, Azuma I, Yamaoka T
2008; 2(3):178-187.

Preliminary research on abating rat testicle toxicity due to triptolide after oral polymer nanoparticle delivery.

Liu MX, Dong J, Yang YJ, Yang XL, Xu HB
2008; 2(3):188-193.

Questionnaire on the awareness of generic drugs among outpatients and medical staff.

Hoshi S, Kimura H
2008; 2(3):194-199.

Transdermal patch incorporating salbutamol sulphate: *In vitro* and clinical characterization.

El-Gendy NA, Sabry NA, El-Attar M, Omar E, Mahmoud M
2008; 2(4):219-228.

Establishment of a cell-based assay to screen insulin-like hypoglycemic drugs.

Zhang L, Hu JJ, Du GH
2008; 2(4):229-233.

Inhibition of *in vivo* angiogenesis by *Anacardium occidentale* L. involves repression of the cytokine VEGF gene expression.

Lingaraju SM, Keshavaiah K, Salimath BP
2008; 2(4):234-244.

Production of a human antibody fragment against the insulin-like growth factor I receptor as a fusion protein.

Kusada Y, Morizono T, Sakai K, Takayanagi A, Shimizu N, Fujita-Yamaguchi Y
2008; 2(4):245-253.

The effects of dietary obesity on protein expressions of insulin signaling pathway in rat aorta.

Fatani SH, Naderali EK, Panchiani S, Shah F, Wong C
2008; 2(4):254-261.

The neuroprotective effect of antidepressant drug *via* inhibition of TIEG2-MAO B mediated cell death.

Lu D, Johnson C, Johnson S, Tazik S, Ou XM
2008; 2(5):289-295.

Regulation of the nitric oxide synthesis pathway and cytokine balance contributes to the healing action of *Myristica malabarica* against indomethacin-induced gastric ulceration in mice.

Maity B, Banerjee D, Bandyopadhyay SK, Chattopadhyay S
2008; 2(5):296-304.

Part 2. Long term *in vivo/in vitro* evaluation of the Cholecystokinin antagonists: *N*-(5-methyl-3-oxo-1,2-diphenyl-2,3-dihydro-1*H*-pyrazol-4-yl)-*N'*-phenylurea MPP and carboxamide MPM.

Lattmann E, Boonprakob Y, Sattayasai J
2008; 2(6):344-352.

Pioglitazone attenuates tactile allodynia and microglial activation in mice with peripheral nerve injury.

Iwai S, Maeda T, Kiguchi N, Kobayashi Y, Fukazawa Y, Ozaki M, Kishioka S
2008; 2(6):353-356.

Phenolphthalein treatment in pregnant women and congenital abnormalities in their offspring: A population-based case-control study.

Báñhidy F, Ács N, Puhó EH, Czeizel AE
2008; 2(6):357-367.